arXiv:1609.08467v1 [math.GR] 27 Sep 2016

A FAMILY OF PERMUTATION GROUPS WITH EXPONENTIALLY MANY NON-CONJUGATED REGULAR ELEMENTARY ABELIAN SUBGROUPS

SERGEI EVDOKIMOV, MIKHAIL MUZYCHUK, AND ILIA PONOMARENKO

Abstract. Given a prime p, we construct a permutation group containing at least p^{p-2} non-conjugated regular elementary abelian subgroups of order p^3. This gives the first example of a permutation group with exponentially many non-conjugated regular subgroups.

1. Introduction

Regular subgroups of permutation groups arise in many natural contexts, for example, in group factorizations [4], Schur rings [6], Cayley graphs [1], etc. In the present paper, given a group H and a permutation group Γ, we are interested in the number of Γ-orbits

$$b_H(\Gamma) := |\operatorname{Orb}(\Gamma, \operatorname{Reg}(\Gamma, H))| \tag{1}$$

in the action of Γ by conjugation on the set $\operatorname{Reg}(\Gamma, H)$ of all its regular subgroups isomorphic to H. Using terminology and arguments of [1], one can see that if Γ is the automorphism group of an object of a concrete category $\mathscr{C}$, then $b_H(\Gamma)$ equals the number of pairwise non-equivalent representations of this object as a Cayley object over H in $\mathscr{C}$. Note that H is a CI-group with respect to the category $\mathscr{C}$ if and only if $b_H(\Gamma) = 1$ for every group $\Gamma = \operatorname{Aut}(X)$, where X is a Cayley object over H in $\mathscr{C}$. As $\mathscr{C}$, one can take, for example, the category of finite graphs or other combinatorial structures.

Let H be a cyclic group. Then, obviously, $b_H(\Gamma) \le c(\Gamma)$, where $c(\Gamma)$ is the number of the conjugacy classes of full cycles contained in Γ. It was proved in [5] that the latter number does not exceed $n = |H|$.[1] Thus, in this case $b_H(\Gamma) \le n$.

The simplest non-cyclic case appears when H is an elementary abelian group E_{p^2}. Here, $b_H(\Gamma) \le b_H(P)$ by the Sylow theorem, where P is a Sylow p-subgroup of the group Γ. To estimate $b_H(P)$, without loss of generality one can assume that P is a transitive p-group of degree p^2, the action of which on some imprimitivity system induces a regular (cyclic) group of order p, i.e., P belongs to the class defined in the same way as the class $\mathcal{E}_p$ in Theorem 1.1 below with p^3 replaced by p^2. With the help of the technique developed in Section 2, one can describe the set $\operatorname{Reg}(\Gamma, H)$ (cf. Theorem 2.2 and Lemma 2.4). Then applying [2, Theorem 6.1], one can prove that $b_H(P) \le p$. Thus, in this case $b_H(\Gamma) \le n$ too.

[1] More exactly, under the Classification of Finite Simple Groups, $c(\Gamma) \le \varphi(n)$ where φ is the Euler function, ibid.

In the above two cases, the number $b_H(\Gamma)$ does not exceed n for all Γ. The main result of the present paper (Theorem 1.1) shows that in the general case, neither this bound nor even substantially weaker bounds are valid.

Theorem 1.1. *Let $H = E_{p^3}$, where p is a prime. Denote by $\mathcal{E}_p$ the class of all transitive p-groups of degree p^3, the action of which on some imprimitivity system induces a regular group isomorphic to E_{p^2}. Then there exists a group $\Gamma \in \mathcal{E}_p$ such that $b_H(\Gamma) \geq p^{p-2}$.*

From Theorem 1.1, it follows that there is no function f, for which the inequality $b_H(\Gamma) \leq n^{f(r)}$ holds for all abelian groups H of rank at most r and all permutation groups Γ of degree n. It would be interesting to find an invariant $t = t(\Gamma)$ such that

$$b_H(\Gamma) \leq n^{f(r,t)}$$

for a function f in r and t; for instance, one can try to take $t = t(\Gamma)$ to be the minimal positive integer t', for which the group Γ is t'-closed as a permutation group in the sense of [7].[2]

The proof of Theorem 1.1 is given in Section 3. It is based on a representation of the groups belonging to the class $\mathcal{E}_p$ with the help of two-variable polynomials over the field $\mathbb{F}_p$. The details are presented in Section 2. It is interesting to note that for every group Γ from Theorem 1.1, the stabilizer of the imprimitivity system is, up to language, a Generalized Reed-Muller code [3].

Notation. As usual, $\mathbb{F}_p$ and $\mathrm{Sym}(V)$ denote the field of order p and the symmetric group on the set V. An elementary abelian p-group of order p^n is denoted by E_{p^n}.

2. Permutation groups and polynomials

Let p be a prime. Denote by R_p the factor ring of the polynomial ring $\mathbb{F}_p[X, Y]$ modulo the ideal generated by polynomials $X^p - 1$ and $Y^p - 1$. The images of the variables X and Y are denoted by x and y, respectively. Denote by V the disjoint union of one-dimensional subspaces

$$V_{i,j} = \{\alpha x^i y^j : \ \alpha \in \mathbb{F}_p\}, \quad i, j = 0, \ldots, p-1,$$

of the ring R_p considered as a linear space over $\mathbb{F}_p$.

Every element $f = \sum_{i,j} \alpha_{i,j} x^i y^j$ of R_p yields a permutation

$$\sigma_f : \alpha x^i y^j \mapsto (\alpha + \alpha_{i,j}) x^i y^j$$

of the set V. This produces a permutation group on V with p^2 orbits $V_{i,j}$ that is isomorphic to the additive group of the ring R_p. For a subgroup I of the latter group, the corresponding subgroup of $\mathrm{Sym}(V)$ is denoted by $\Delta(I)$. In addition, we define two commuting permutations

$$\tau_x : \alpha x^i y^j \mapsto \alpha x^{i+1} y^j, \qquad \tau_y : \alpha x^i y^j \mapsto \alpha x^i y^{j+1}.$$

Clearly, each of them commutes with the permutation $s = \sigma_{f_0}$, where $f_0 = \sum_{i,j} x^i y^j$. The following statement is straightforward.

Lemma 2.1. *In the above notation, we have*

(1) $\tau_x^{-1} \sigma_f \tau_x = \sigma_{fx}$ *and* $\tau_y^{-1} \sigma_f \tau_y = \sigma_{fy}$ *for all* $f \in R_p$,

[2]Here for groups $\Gamma \in \mathcal{E}_p$, the upper bound in inequality (8) could be useful.

(2) $G_0 := \langle s, \tau_x, \tau_y \rangle$ *is a regular group on* V *isomorphic to* E_{p^3}.

Set $\Gamma(I)$ to be the group generated by $\Delta(I)$ and τ_x, τ_y. If I is an ideal of R_p, then in view of statement (1) of Lemma 2.1,

$$\Delta(I) \trianglelefteq \Gamma(I) \quad \text{and} \quad \Gamma(I)/\Delta(I) \cong E_{p^2}. \tag{2}$$

If I is not an ideal, then $\Gamma(I) = \Gamma(I')$, where I' is the ideal of R_p generated by I.

Theorem 2.2. *Let* p *be a prime. Then*

(1) *for every ideal* $I \neq 0$ *of the ring* R_p*, the group* $\Gamma(I)$ *belongs to the class* $\mathcal{E}_p$*,*
(2) *every group* $\Gamma \in \mathcal{E}_p$ *with* $b_H(\Gamma) > 0$ *is permutation isomorphic to the group* $\Gamma(I)$ *for some ideal* I *of* R_p*.*

Proof. To prove statement (1), let $I \neq 0$ be an ideal of R_p. Then at least one of the sets $V_{i,j}$ is an orbit of the group $\Delta(I)$. Since τ_x and τ_y commute, the group $\langle \tau_x, \tau_y \rangle$ acts regularly on the set $S = \{V_{i,j} : \ i, j = 0, \dots, p-1\}$. This implies that the group $\Gamma(I)$ is transitive and S is an imprimitivity system of it. The action of $\Gamma(I)$ on this system induces a regular group isomorphic to E_{p^2} that is generated by the images of τ_x and τ_y with respect to this action. Thus, $\Gamma(I) \in \mathcal{E}_p$.

Let $\Gamma \in \mathcal{E}_p$. Then Γ is a transitive p-group of degree p^3, the action of which on some imprimitivity system S' induces a regular group isomorphic to E_{p^2}. Without loss of generality, we may assume that $\Gamma \leq \mathrm{Sym}(V)$ with V as above. Furthermore, since $b_H(\Gamma) > 0$, the group Γ contains a regular subgroup G' isomorphic to $H = E_{p^3}$. Choose an element $s' \in G'$ such that $\mathrm{Orb}(\langle s' \rangle, V) = S'$. Then there exists a group isomorphism

$$\varphi : G' \to G_0$$

taking s' to s (see statement (2) of Lemma 2.1). Since φ is induced by a permutation of V, we may assume that $S' = S$ and $G_0 \in \mathrm{Reg}(\Gamma, E_{p^3})$. Note that the permutation s belongs to the stabilizer Δ of the blocks $V_{i,j}$ in Γ. Therefore, $\mathrm{Orb}(\Delta, V) = S$. Since the restriction of Δ to $V_{i,j}$ is a p-group of degree p that contains the restriction of s to $V_{i,j}$ for all i, j, this implies that

$$\Delta \leq \Delta(R_p).$$

It follows that $\Delta = \Delta(I)$ for a subgroup I of R_p. Taking into account that Δ is normalized by τ_x and τ_y, we conclude that I is an ideal of R_p by statement (1) of Lemma 2.1. □

Any maximal element in the class $\mathcal{E}_p$ is permutation isomorphic to the (imprimitive) wreath product of regular groups isomorphic to E_p and E_{p^2}. One of these maximal elements equals the group $\Gamma_p := \Gamma(R_p)$; set also $\Delta_p = \Delta(R_p)$. We need two auxiliary lemmas.

Lemma 2.3. *Let* $g, h \in R_p$*. Then the order of the permutation* $t_{g,x} = \sigma_g \tau_x$ *(resp.* $t_{h,y} = \sigma_h \tau_y$*) equals* p *if and only if* $g \in aR_p$ *(resp.* $h \in bR_p$*), where* $a = x - 1$ *and* $b = y - 1$*.*

Proof. Let $g = \sum_{i,j} \alpha_{i,j} x^i y^j$, and let $v = \alpha x^i v^j$ be a point of V. Then by the definition of $t_{g,x}$, we have

$$v^{t_{g,x}} = (\alpha + \alpha_{i,j}) x^{i+1} y^j.$$

This implies that the order of $t_{g,x}$ equals p if and only if the following condition is satisfied:

$$\sum_{i=0}^{p-1} \alpha_{i,j} = 0, \qquad j = 0, \dots, p-1. \tag{3}$$

Note that this is always true, whenever $g \in aR_p$. Conversely, suppose that relations (3) hold for some $g \in R_p$. Then

$$\alpha_{0,j} = \alpha'_{1,j} - \alpha'_{0,j}, \ \dots, \ \alpha_{p-1,j} = \alpha'_{0,j} - \alpha'_{p-1,j},$$

where $\alpha'_{i,j} = \sum_{k=0}^{i-1} \alpha_{k,j}$ for all i, j. It follows that $g = ag'$ with $g' = \sum_{i,j} \alpha'_{i,j} x^i y^j$. This completes the proof of the first statement. The second statement (on the order of $t_{h,y}$) is proved similarly. □

Lemma 2.4. *A permutation group G belongs to the set $\mathrm{Reg}(\Gamma_p, E_{p^3})$ if and only if there exist elements $g \in aR_p$ and $h \in bR_p$ such that*

$$G = \langle s, t_{g,x}, t_{h,y} \rangle \quad \textit{and} \quad ah = bg. \tag{4}$$

Proof. To prove the "only if" part, suppose that $G \in \mathrm{Reg}(\Gamma_p, E_{p^3})$. Then G is a self-centralizing subgroup of $\mathrm{Sym}(V)$. On the other hand, the centralizer of G in $\mathrm{Sym}(V)$ contains the central element s of the group Γ_p. Thus, $s \in G$. The other two generators of G can obviously be chosen so that their images with respect to the epimorphism $\Gamma_p \to \Gamma_p/\Delta_p$ coincide with x and y. By Lemma 2.3, this implies that there exist $g \in aR_p$ and $h \in bR_p$, for which the first equality in (4) holds. Next, since the group G is abelian, the definition of $t_{g,x}$ and $t_{h,y}$ implies that

$$\sigma_g \tau_x \, \sigma_h \tau_y = t_{g,x} t_{h,y} = t_{h,y} t_{g,x} = \sigma_h \tau_y \, \sigma_g \tau_x.$$

Each of the permutations on the left- and right-hand sides takes the point $\alpha x^i y^j \in V_{i,j}$ to a certain point $\alpha' x^{i+1} y^{j+1} \in V_{i+1,j+1}$. Calculating the images of the former point with respect to them, we obtain

$$\alpha + g_{i,j} + h_{i+1,j} = \alpha' = \alpha + h_{i,j} + g_{i,j+1}$$

or, equivalently, $h_{i+1,j} - h_{i,j} = g_{i,j+1} - g_{i,j}$ for all i, j. Therefore, $ah = xh - h = yg - g = bg$, as required.

Conversely, let G be the group defined by relations (4). Then the above argument shows that the permutations s, $t_{g,x}$, and $t_{h,y}$ pairwise commute. Therefore, the group G is abelian. Moreover, the definition of s and Lemma 2.3 imply that G is elementary abelian and transitive. Thus, $G \in \mathrm{Reg}(\Gamma_p, E_{p^3})$, as required. □

3. Proof of Theorem 1.1

By statement (1) of Theorem 2.2, we may restrict ourselves to looking for the group Γ of the form $\Gamma(I)$, where I is an ideal of the ring R_p.

For every integer $k \geq 0$, set

$$I_k = \mathrm{span}_{\mathbb{F}_p}\{a^i b^j : \ i + j \geq k\},$$

where the elements a and b are as in Lemma 2.4. Clearly, I_k is an ideal of R_p, and $I_{k+1} \subseteq I_k$ for all k, and also $I_k = 0$ for $k > 2(p-1)$. Below, the kernels of the mappings $I_k \to aI_k$ and $I_k \to bI_k$ induced by the multiplication by a and b are denoted by A_k and B_k, respectively.

Lemma 3.1. *Suppose that $p \le k \le 2(p-1)$. Then*

(1) $\dim(I_k) = \binom{2p-k}{2}$,
(2) $aI_k = bI_k = I_{k+1}$,
(3) $\dim(A_k) = \dim(B_k) = 2p-k-1$.

Proof. The leading monomials of the polynomials

$$(x-1)^i(y-1)^j, \qquad 0 \le i,j \le p-1,$$

are obviously linearly independent. Therefore, the polynomials a^ib^j with $i+j \ge k$ form a linear basis of the ideal I_k. This immediately proves statement (1). To prove statement (2), we note that, obviously, $aI_k \subseteq I_{k+1}$. Conversely, let $c \in I_{k+1}$. Since $k \ge p$, we have $c = abu$ for some $u \in I_{k-1}$, which proves the reverse inclusion. The rest of statement (2) is proved similarly. Finally, statement (3) follows, because the linear space A_k (resp. B_k) is spanned by the monomials $a^{p-1}b^i$ (resp. a^ib^{p-1}) with $k-p+1 \le i \le p-1$. □

In what follows, for a subgroup G of a group Γ we denote by G^Γ the set of all Γ-conjugates of G.

Lemma 3.2. *Let $\Gamma_{k,p} = \Gamma(I_k)$, where k is as in Lemma 3.1. Then*

(1) $|\Gamma_{k,p}| = p^{2+\dim(I_k)}$,
(2) $|\operatorname{Reg}(\Gamma_{k,p}, E_{p^3})| = p^{\dim(A_k)+\dim(B_k)+\dim(I_{k+1})-2}$,
(3) $p^{\dim(I_k)-4} \le |G^{\Gamma_{k,p}}| \le p^{\dim(I_k)-1}$ *for all* $G \in \operatorname{Reg}(\Gamma_{k,p}, E_{p^3})$.

Proof. Obviously, $|\Delta(I_k)| = p^{\dim(I_k)}$. Therefore, statement (1) follows from the right-hand side of formula (2). Next, from Lemma 2.4 it follows that

$$\operatorname{Reg}(\Gamma_{k,p}, E_{p^3}) = \{G_{g,h} : \ (g,h) \in M\},$$

where $G_{g,h} = \langle s, t_{g,x}, t_{h,y}\rangle$ and

$$M = \{(g,h) \in (I_k \cap aR_p) \times (I_k \cap bR_p) : \ ah = bg\}. \tag{5}$$

However, $I_k \cap aR_p = I_k \cap bR_p = I_k$, because $k \ge p$. So by statement (2) of Lemma 3.1, the element $ah = bg$ runs over the ideal I_{k+1}, whenever (g,h) runs over the set M. By the definition of A_k and B_k, this implies that

$$|M| = p^{\dim(A_k)+\dim(B_k)+\dim(I_{k+1})}.$$

Thus to complete the proof of statement (2), it suffices to verify that $G_{g,h} = G_{g',h'}$ if and only if $t_{g,x} = s^i t_{g',x}$ and $t_{h,y} = s^j t_{h',y}$ for some $0 \le i,j \le p-1$. However, this is true, because $G_{g,h} = G_{g',h'}$ if and only if $\varphi(G_{g,h}) = \varphi(G_{g',h'})$, where φ is the quotient epimorphism of $\Gamma_{k,p}$ modulo the group $\langle s\rangle$.

To prove statement (3), we note that in view of statement (1),

$$|G^\Gamma| = \frac{|\Gamma|}{|N|} = \frac{p^{2+\dim(I_k)}}{|C| \cdot |N/C|}, \tag{6}$$

where $\Gamma = \Gamma_{k,p}$, and N and C are, respectively, the normalizer and centralizer of G in Γ. Since G is a regular elementary abelian group and the quotient N/C is isomorphic to a subgroup of a Sylow p-subgroup P of the group $\operatorname{Aut}(G) \cong \mathrm{GL}(3,p)$ (here we use the fact that Γ is a p-group), we conclude that

$$|C| = |G| = p^3 \quad \text{and} \quad 1 \le |N/C| \le |P|.$$

However, $|P| = p^3$. Thus, statement (3) follows from formula (6). □

To complete the proof of Theorem 1.1, we note that $\mathrm{Reg}(\Gamma_k, E_{p^3})$ is the disjoint union of distinct sets G^{Γ_k}, where $\Gamma_k = \Gamma_{k,p}$ as in Lemma 3.2 and $G \in \mathrm{Reg}(\Gamma_k, E_{p^3})$. Therefore, setting m_k and M_k to be, respectively, the minimum and maximum of the numbers $|G^{\Gamma_k}|$, we obtain

$$\frac{|\mathrm{Reg}(\Gamma_k, E_{p^3})|}{m_k} \geq b_H(\Gamma_k) \geq \frac{|\mathrm{Reg}(\Gamma_k, E_{p^3})|}{M_k}. \tag{7}$$

However, by statement (3) of Lemma 3.2, $m_k \geq p^{\dim(I_k)-4}$ and $M_k \leq p^{\dim(I_k)-1}$. By statement (2) of Lemma 3.2, this implies that

$$\frac{|\mathrm{Reg}(\Gamma_k, E_{p^3})|}{m_k} \leq p^{d+2} \quad \text{and} \quad \frac{|\mathrm{Reg}(\Gamma_k, E_{p^3})|}{M_k} \geq p^{d-1}.$$

where $d = \dim(A_k) + \dim(B_k) + \dim(I_{k+1}) - \dim I_k$. Besides, by statements (1) and (3) of Lemma 3.1, we have $d = 2p - k - 1$. Thus,

$$p^{2p-k+1} \geq b_H(\Gamma_k) \geq p^{2p-k-2}. \tag{8}$$

This lower bound for $b_H(\Gamma_k)$ with $k = p - 1$ proves Theorem 1.1.

References

[1] L. Babai, *Isomorphism problem for a class of point symmetric structures*, Acta Math. Acad. Sci. Hung., **29**, No. 3, 329–336 (1977).

[2] S. Evdokimov and I. Ponomarenko, *Recognizing and isomorphism testing circulant graphs in polynomial time*, Algebra and Analysis, **15**, No. 6, 1–34 (2003).

[3] T. Kasami, S. Lin, and W. W. Peterson, *New generalisations of the Reed-Muller codes*, EEE Trans. Information Theory, **IT-14**, 189–199 (1968).

[4] M. W. Liebeck, C. Praeger, and J. Saxl, *Regular subgroups of primitive permutation groups*, Memoirs Amer. Math. Soc., **203**, No. 952, 1–88 (2010).

[5] M. Muzychuk, *On the isomorphism problem for cyclic combinatorial objects*, Discr. Math., **197/198**, 589–606 (1999).

[6] H. Wielandt, *Finite permutation groups*, Academic press, New York - London (1964).

[7] H. Wielandt, *Permutation groups through invariant relations and invariant functions*, Lect. Notes Dept. Math. Ohio St. Univ., Columbus (1969).

Steklov Institute of Mathematics at St. Petersburg, Russia
E-mail address: evdokim@pdmi.ras.ru

Netanya Academic College, Netanya, Israel
E-mail address: muzy@netanya.ac.il

Steklov Institute of Mathematics at St. Petersburg, Russia
E-mail address: inp@pdmi.ras.ru